Stanford University Library
JUL 25 1906

JULY 1906

ANNALS OF MATHEMATICS

(FOUNDED BY ORMOND STONE)

EDITED BY

ORMOND STONE W. E. BYERLY F. S. WOODS
MAXIME BÔCHER E. V. HUNTINGTON

PUBLISHED UNDER THE AUSPICES OF HARVARD UNIVERSITY

SECOND SERIES Vol. 7 No. 4

FOR SALE BY
THE PUBLICATION OFFICE OF HARVARD UNIVERSITY
2 University Hall, Cambridge, Mass., U. S. A.

London: LONGMANS, GREEN & Co.
39 Paternoster Row

Leipzig: OTTO HARRASSOWITZ
Querstrasse 14

The following pamphlets, most of which have been reprinted from the ANNALS OF MATHEMATICS, second series, will be sent post-paid on receipt of price by the PUBLICATION OFFICE OF HARVARD UNIVERSITY, 2 University Hall, Cambridge, Mass.

MAXIME BÔCHER, Regular Points of Linear Differential Equations of the Second Order. 23 pp. 1896. $0.50

W. F. OSGOOD, Introduction to Infinite Series. 71 pp. Second Edition, 1902. .75

JAMES PIERPONT, Galois' Theory of Algebraic Equations. 67 pp. 1900. *From vols.* 1–2. .75

MAXIME BÔCHER, The Theory of Linear Dependence. 16 pp. 1901. *From vol.* 2. .35

W. F. OSGOOD, Sufficient Conditions in the Calculus of Variations. 25 pp. 1901. *From vol.* 2. .50

F. S. WOODS, Space of Constant Curvature. 42 pp. 1902. *From vol.* 3. .50

W. F. OSGOOD, The Integral as the Limit of a Sum, and a Theorem of Duhamel's. 18 pp. 1903. *From vol.* 4. .35

E. V. HUNTINGTON, The Continuum as a Type of Order: An Exposition of the Modern Theory. With an Appendix on the Transfinite Numbers. 63 pp 1905. *From vols.* 6–7. .50

MAXIME BÔCHER, Introduction to the Theory of Fourier's Series. 72 pp. 1906. *From vol.* 7. .75

A PROOF OF THE THEOREM CONCERNING ARTIFICIAL SINGULARITIES

By D. R. Curtiss

In the January number of the *Bulletin of the American Mathematical Society*,* Professor Landau has supplied data lacking in certain fallacious proofs of the well-known theorem which states that if $f(z)$ is single-valued and analytic at all points of the neighborhood of the point c, exclusive of c, and remains finite throughout this neighborhood, it has at most an artificial singularity at c; i. e., with a suitable definition of $f(c)$ the function $f(z)$ will be analytic at c.†

I wish to point out in this note a simple proof of this theorem which does not assume, as does Professor Landau's paper, the existence of $\lim_{z=c} f(z)$.

The proof makes use of an auxiliary function $\psi(z)$ defined by the equations

$$\psi(z) = (z-c)^2 f(z) \qquad (z \neq c), \qquad \psi(c) = 0.‡$$

We have

$$\psi'(z) = 2(z-c)f(z) + (z-c)^2 f'(z) \qquad (z \neq c),$$
$$\psi'(c) = 0.$$

Thus $\psi(z)$ has a derivative at c as well as throughout its neighborhood. Though Goursat's theorem § makes it unnecessary to prove $\psi'(z)$ continuous

---

* Vol. 12, pp. 155–156.

† For references and a discussion of various proofs of this theorem see Professor Osgood's paper, "Some points in the elements of the theory of functions," *Bull. Amer. Math. Soc.*, vol. 2 (1896), pp. 296–302.

‡ The auxiliary function considered by Professor Landau and used in the above-mentioned incorrect proofs was the function $\phi(z)$ defined as follows:

$$\phi(z) = (z-c)f(z) \qquad (z \neq c), \qquad \phi(c) = 0.$$

§ *Trans. Amer. Math. Society*, vol. 1 (1900), pp. 14–16.

at c, we may note that this property is easily deduced from the formula, valid when $|z-c|$ is sufficiently small,

$$f''(z) = \frac{1}{2\pi i}\int_C \frac{f(t)\,dt}{(t-z)^2},$$

where C is a circumference about z of radius $\frac{1}{2}|z-c|$. If $M > |f(z)|$ throughout the neighborhood of c, exclusive of c, we have

$$|f''(z)| < \frac{2M}{|z-c|};$$

hence

$$\lim_{z=c} \psi'(z) = \psi'(c) = 0.$$

But it can be shown, either by the use of Taylor's series or directly from Cauchy's formula written in the form

$$\begin{aligned}\psi(z) &= \frac{1}{2\pi i}\int \frac{\psi(t)\,dt}{t-z} \\ &= \frac{1}{2\pi i}\int \frac{\psi(t)\,dt}{t-c} + \frac{z-c}{2\pi i}\int \frac{\psi(t)\,dt}{(t-c)^2} + \frac{(z-c)^2}{2\pi i}\int \frac{\psi(t)\,dt}{(t-z)(t-c)^2} \\ &= \psi(c) + (z-c)\,\psi'(c) + (z-c)^2\,\Psi(z),^*\end{aligned}$$

that on account of the relation $\psi(c) = \psi'(c) = 0$, $\psi(z)$ can be expressed as the product of $(z-c)^2$ and a function $\Psi(z)$ analytic at the point c as well as throughout its neighborhood. We have therefore only to add the definition $f(c) = \Psi(c)$ to make $f(z)$ identical with $\Psi(z)$ throughout the neighborhood of c, inclusive of c.

EVANSTON, ILLINOIS,
JANUARY, 1906.

* The corresponding general form of Taylor's series with the remainder, although capable of wide application, has received little notice in treatises on the theory of functions.

ANOTHER PROOF OF THE THEOREM CONCERNING ARTIFICIAL SINGULARITIES

By Maxime Bôcher

The theorem considered by Professor Curtiss in the preceeding note is so important that still another simple proof may not be out of place. The proof which I give is a simplification of the second proof given by Osgood in the *Bulletin of the American Mathematical Society* for June 1896, p. 301.

Lemma. *If $f(z)$ is continuous throughout a region S, and is analytic at every point of this region except at a point c which is not on the boundary of S, then f is analytic at c also.*

Consider a circle K with c as centre and lying wholly in S. The integral of $f(z)$ taken around any regular closed path in K is zero. To prove this we consider three cases:

1) If this closed path does not surround or pass through c, $f(z)$ is analytic everywhere upon and within the path. Hence the integral is zero.

2) If the path surrounds c, the integral around it is equal to the integral around an arbitrarily small circle having c as centre. Accordingly its absolute value cannot exceed $2\pi rM$ where r is the radius of this small circle and M is the maximum of $|f(z)|$ on its circumference. But since as r approaches zero, $\lim M = |f(c)|$, we have $\lim 2\pi rM = 0$. Hence the value of our integral must be zero.

3) If the path passes through c, the integral around it is equal to the integral along an arbitrarily short arc of length l of this path having c as its middle point plus the integral along a path connecting the extremities of this arc and of length less than $2l$. Accordingly the absolute value of this integral cannot exceed $3lM$, where M is the maximum of $|f(z)|$ on the path just described. Since this vanishes with l, our integral is again zero.

Since $f(z)$ is continuous throughout K and its integral around any regular closed path in K is zero, it follows from Morera's theorem * that $f(z)$ is analytic at c as well as elsewhere.

THEOREM. *If $f(z)$ is analytic throughout a region S except at a point c which is not on the boundary of S, and if $f(z)$ remains finite in the neighborhood of c, then $f(z)$ approaches a finite limit as z approaches c, and if this limiting value be taken as the definition of $f(c)$, then f is analytic at c.*

For the function $(z - c)f(z)$ is by our lemma analytic at c if we define it as having the value zero there. Hence we may write

$$(z - c)f(z) = a_1(z - c) + a_2(z - c)^2 + \cdots,$$

and therefore, when $z \neq c$,

$$f(z) = a_1 + a_2(z - c) + \cdots.$$

Hence as z approaches c, $\lim f(z) = a_1$, and if we define $f(c) = a_1$, f is analytic at c.

HARVARD UNIVERSITY,
CAMBRIDGE, MASS.

* Cf. Osgood, *Funktionentheorie*, vol. 1, p. 256. This theorem is so easily proved and so useful that it may well be introduced at an early stage even in an elementary course on the theory of functions.

CURVES OF MINIMUM MOMENT OF INERTIA WITH RESPECT TO A POINT

By Max Mason

The problem to be considered in the following pages furnishes an example of the calculus of variations in which discontinuous solutions play an important part. The problem may be stated as follows:

Among the curves that join two given points, 1 *and* 2, *that one is required which has the least moment of inertia with respect to a third point* 0.*

It is evident that only those curves which lie in the plane 012 need be considered. In fact, the orthogonal projection, C', on this plane of any curve C not lying in the plane has a smaller moment of inertia than the curve C, since C is longer than C' and each point of C is farther from 0 than is the corresponding point of C'.

Let the polar coordinates of the points 1, 2 be (r_1, θ_1), (r_2, θ_2), the point 0 being chosen as the pole. Assuming the equation of the curve in parameter representation,

$$r = r(t), \ \theta = \theta(t),$$

it is required to minimize the integral

$$J = \int r^2 ds = \int_{t_1}^{t_2} r^2 \sqrt{r'^2 + r^2\theta'^2}\, dt,$$

under the conditions

$$r(t_1) = r_1, \quad r(t_2) = r_2, \quad \theta(t_1) = \theta_1, \quad \theta(t_2) = \theta_2.$$

The equation of the extremals† may be found most simply by writing the integrand of J in the form

$$r^2 \sqrt{1 + r^2 \left(\frac{d\theta}{dr}\right)^2}\, dr.$$

---

* The density is supposed constant and the same for all curves. The problem has not been previously treated beyond the point of determining the equation of the extremals. This was done by Bonnet, *Journal de Mathématiques*, vol. 9 (1844), p. 97.

† For the notation and well known theorems of the calculus of variations the reader is referred to Bolza, *Lectures on the Calculus of Variations*, Chicago, 1904.

Since this does not contain θ we may write down at once a first integral of the differential equation of the extremals, namely,

$$\frac{r^4 \frac{d\theta}{dr}}{\sqrt{1 + r^2\left(\frac{d\theta}{dr}\right)^2}} = \text{constant} = \beta,$$

or,

$$\frac{d\theta}{dr} = \frac{\beta}{r\sqrt{r^6 - \beta^2}}.$$

For $\beta = 0$ the extremals are straight lines through the pole, $\theta = \text{const}$. For general values of β the integration of the above equation gives

$$3\theta + \alpha = \arctan \frac{\sqrt{r^6 - \beta}}{\beta},$$

or,

$$\beta r^3 = \sec (3\theta + \alpha), \tag{1}$$

where α is a second constant of integration. An extremal $\theta = \text{const}$ will be denoted by S, the remaining extremals by C.

1. The straight line solutions. In case the three points 0, 1, 2 lie in a straight line the problem may be solved without reference to the general theory. The extremal joining 1 and 2 is the straight line S through 0, 1, 2. Suppose first that the points 1 and 2 are not separated by 0. Let the order be 0, 1, 2. Denote by $\bar{C}$ any curve other than S which joins 1 and 2, and by s the length of arc on $\bar{C}$ or S measured from 2. Now

$$J = \int r^2 ds;$$

the length of $\bar{C}$ is greater than that of S, and the value of r on $\bar{C}$ is greater than the value of r on S for the same value of s. Hence $J(\bar{C}) > J(S)$. If the points 1 and 2 are separated by 0 there is a point 3 on S such that the lengths of S_{23} and S_{13} are greater than those of $\bar{C}_{20}$ and $\bar{C}_{10}$ respectively. Then the above inequality holds for the same reason as before.

We may therefore state the following theorem:

THEOREM I. *If the three points* 0, 1, 2 *lie on a straight line, the segment of that line between* 1 *and* 2 *has the least moment of inertia with respect to* 0 *of all curves joining* 1 *and* 2.

2. The Continuous Solutions. It is evident from equation (1) that unless $|\theta_2 - \theta_1| < \pi/3$ the points 1 and 2 cannot be joined by an extremal. We will assume for the present that this inequality holds. Since the extremal of the set (1) corresponding to the values α, β is identical with that for the values $\alpha + \pi$, $-\beta$ we may assume without restriction that

$$0 \leqq \alpha < \pi.$$

To determine an extremal joining 1 and 2, the equations

$$\beta r_1^3 = \sec(3\theta_1 + \alpha)$$
$$\beta r_2^3 = \sec(3\theta_2 + \alpha)$$

must be solved for α and β. The value of β is uniquely determined when α is known. Eliminating β we have

$$r_2^3(\cos 3\theta_2 \cos\alpha - \sin 3\theta_2 \sin\alpha) = r_1^3(\cos 3\theta_1 \cos\alpha - \sin 3\theta_1 \sin\alpha),$$

or,

$$\tan\alpha = \frac{r_1^3 \cos 3\theta_1 - r_2^3 \cos 3\theta_2}{r_1^3 \sin 3\theta_1 - r_2^3 \sin 3\theta_2}.$$

Since $0 \leqq \alpha < \pi$ this equation admits a unique solution. Hence:

If $|\theta_2 - \theta_1| < \pi/3$ the points $1(r_1, \theta_1)$, $2(r_2, \theta_2)$ may be connected by one and only one extremal C.

Let α_0, β_0 be the values of α, β which determine the extremal C. If we allow β to vary we obtain a one-parameter family of extremals, all having the same pair of radii vectores as asymptotes. These extremals form a field. In fact, the set being given by the equation

$$r^3 = \frac{\sec(3\theta + \alpha_0)}{\beta},$$

we have

$$\frac{\partial r}{\partial \beta} = -\frac{\sec(3\theta + \alpha_0)}{3r^2\beta^2} = -\frac{r}{3\beta},$$

which is different from zero within the angle formed by the asymptotes.

The extremal C may therefore be imbedded in a field which covers the entire interior of the area bounded by the asymptotes to C.

Since the function

$$F_1 = \frac{F_{\theta'\theta'}}{r'^2} = \frac{r^4}{(r'^2 + r^2\theta'^2)^{3/2}}$$

is positive for all directions (r', θ') it follows the Weierstrass E-function, $E(r, \theta, r', \theta', \bar{r}', \bar{\theta}')$, is never negative in the field, and is only then zero, when the direction $(\bar{r}', \bar{\theta}')$ coincides with the direction (r', θ'). *Weierstrass's condition is therefore satisfied.*

The sufficient conditions being thus satisfied the extremal C_{12} minimizes the integral with respect to all curves $\bar{C}_{12}$ which lie within the field.

If the comparison curve $\bar{C}_{12}$ does not lie entirely within the field two cases may arise; the closed curve $\bar{C}_{12} + \bar{C}_{21}$ formed by $\bar{C}_{21}$ and $\bar{C}_{12}$ may or may not include the point 0. If it does not include 0, the curve $\bar{C}_{12}$ may be replaced by a curve C_{12}^* formed by the curves S_{13}, $\bar{C}_{34}$, S_{42} (figure 1) which by theorem



FIG. 1.



FIG. 2.

I gives a smaller value to J than the value $J(\bar{C}_{12})$. But the curve C_{12}^* lies entirely within the field; hence

$$J(C_{12}) < J(C_{12}^*) < J(\bar{C}_{12}).$$

If $\bar{C}_{21} + C_{12}$ does include the point 0 then the line S_{10} if extended will meet $\bar{C}_{12}$ at a point 3, lying at or beyond 0 on S_{10} (figure 2). Then by theorem I:

$$J(S_{10}) + J(S_{03}) < J(\bar{C}_{13}),$$
$$J(S_{02}) < J(S_{03}) + J(\bar{C}_{32}),$$

Hence, adding these inequalities:

$$J(S_{1032}) < J(\bar{C}_{12}).$$

It remains to be proved that

$$J(C_{12}) < J(S_{102}).$$

Let 3, 4 be two points on S_{10} and S_{02} respectively at a distance ϵ from 0. Denote by $C^{\epsilon}_{1\epsilon 2}$ the curve formed by S_{13}, the straight line joining 3 to 4, and S_{42}. Since this curve lies entirely in the field we have

$$J(C^{\epsilon}_{1\epsilon 2}) - J(C_{12}) = \int_{C^{\epsilon}_{1\epsilon 2}} E\,dt,$$

where E is the Weierstrass E function. Hence

$$J(S_{102}) - J(C_{12}) = \lim_{\epsilon=0} \int_{C^{\epsilon}_{1\epsilon 2}} E\,dt > 0.$$

The results may be stated in the following theorem:

THEOREM II. *The extremal*

$$\beta\, r^3 = \sec(3\theta + a)$$

joining the points $1(r_1, \theta_1)$, $2(r_2, \theta_2)$, *where* $|\theta_2 - \theta_1| < \pi/3$, *has the least moment of inertia of all curves joining* 1 *and* 2.

3. The Discontinuous Solutions. If $|\theta_2 - \theta_1| \geqq \pi/3$ the points 1 and 2 can not be joined by an extremal having a continuously turning tangent. Any two points may however be joined in an infinite number of ways by a curve composed of portions of different extremals whose intersections are corners of the curve. A necessary condition that such a curve minimize the integral is that

$$F_1(r, \theta, r', \theta') = \frac{r^4}{(r'^2 + r^2\theta'^2)}$$

shall vanish for some direction (r', θ') when the coordinates of a corner are substituted for r and θ. For the problem under consideration this can only happen at the point 0. The only curve which could form a discontinuous solution is therefore the curve S_{102} composed of the lines $\theta = \theta_1$ and $\theta = \theta_2$ which connect 1 and 2 and meet at 0.

It has been seen above that this curve cannot yield an absolute minimum in case $|\theta_2 - \theta_1| < \pi/3$. Moreover it does not give even a relative minimum to J. In fact, any two points $1'$, $2'$ on S_{10} and S_{02} respectively may be joiued

by an extremal C however near $1'$ and $2'$ may be to 0. As has been shown above

$$J(C_{1'2'}) < J(S_{1'02'}),$$

and hence the curve composed of the segments $S_{11'}$, $C_{1'2'}$, $S_{2'2}$, which may lie as near as we please to S_{102}, gives a smaller value to J than S_{102}. *Hence, if* $|\theta_2 - \theta_1| < \pi/3$, *there exists no discontinuous solution.*

It remains to investigate the existence of discontinuous solutions for the case $|\theta_2 - \theta_1| \geqq \pi/3$.

The value of J taken along an extremal C between two points (r_1, θ_1), (r_2, θ_2) for which $|\theta_2 - \theta_1| < \pi/3$ is

$$J(C_{12}) = \int_{r_0}^{r_1} r^2 \sqrt{1 + r^2 \left(\frac{d\theta}{dr}\right)^2}\, dr + \int_{r_0}^{r_2} r^2 \sqrt{1 + r^2 \left(\frac{d\theta}{dr}\right)^2}\, dr,$$

where r_0 is the minimum value of r on the extremal C_{12}. On substituting the value of $\dfrac{d\theta}{dr}$ from equation (1) this becomes

$$J(C_{12}) = \int_{r_0}^{r_1} \frac{r^5 dr}{\sqrt{r^6 - \beta^{-2}}} + \int_{r_0}^{r_2} \frac{r^5 dr}{\sqrt{r^6 - \beta^{-2}}}.$$

If the points 1 and 2 move so that $|\theta_2 - \theta_1|$ approaches $\pi/3$, then β increases without limit and r_0 approaches 0. Hence

$$\lim_{|\theta_2 - \theta_1| = \pi/3} J(C_{12}) = \frac{r_1^3 + r_2^3}{3}.$$

But

$$J(S_{102}) = \int_0^{r_1} r^2 dr + \int_0^{r_2} r^2 dr = \frac{r_1^3 + r_2^3}{3},$$

and therefore

$$\lim_{|\theta_2 - \theta_1| = \pi/3} \{J(C_{12}) - J(S_{102})\} = 0. \tag{2}$$

We may now compare the value $J(S_{102})$, for the case $|\theta_2 - \theta_1| = \pi/3$, with the value $J(\bar{C}_{12})$ of the integral taken along any other curve $\bar{C}_{12}$ joining 1 and 2. If $\bar{C}$ does not cross the acute angle formed by the lines $\theta = \theta_1$, $\theta = \theta_2$ then $J(S_{102}) < J(\bar{C}_{12})$. This may be easily shown by the method used in proving

theorem I. Suppose that $\bar{C}$ crosses the angle, intersecting S_{10} at 3 and S_{02} at 4 (see figure 3). By theorem I,

$$J(\bar{C}_{13}) + J(\bar{C}_{42}) \geqq J(S_{13}) + J(S_{42}), \tag{3}$$

the equality holding only when $\bar{C}$ coincides with S_{10} from 1 to 3 and with S_{02} from 4 to 2. Let 3′ and 4′ be two points on $\bar{C}$ near 3 and 4 respectively



FIG. 3.

and within the angle formed by S_{102}. Then 3′ and 4′ may be joined by an extremal C, and

$$J(\bar{C}_{3'4'}) - J(C_{3'4'}) = \int_{\bar{C}_{3'4'}} E dt.$$

Hence, by equation (2):

$$J(\bar{C}_{34}) - J(S_{304}) = \lim_{3'=3,\, 4'=4} \int_{\bar{C}_{3'4'}} E\, dt > 0.$$

Adding this inequality to (3) we have

$$J(\bar{C}_{12}) > J(S_{102}).$$

The curve S_{102} therefore minimizes the integral with respect to all other curves joining 1 and 2, if $|\theta_2 - \theta_1| = \pi/3$.

Suppose now that the end points 1, 2 are so situated that $|\theta_2 - \theta_1| > \pi/3$. It will be shown that the curve S_{102} is the solution in this case.

Let C_{12} be any other curve connecting 1 and 2, and $3(r_3, \theta_3)$ a point on $\bar{C}_{12}$ such that $|\theta_3 - \theta_1| = \pi/3$. From the preceding case we have

$$J(\bar{C}_{12}) > J(S_{103}) + J(\bar{C}_{32}),$$

or,

$$J(\bar{C}_{12}) > \frac{r_1^3 + r_3^3}{3} + J(\bar{C}_{32}).$$

Now

$$J(\bar{C}_{32}) = \int_3^2 r^2 \sqrt{r'^2 + r^2\theta'^2}\, dt \geqq \int_3^2 r^2 |r'|\, dt,$$

$$J(\bar{C}_{32}) \geqq \left| \int_3^2 r^2 r' dt \right| = \frac{|r_2^3 - r_3^3|}{3} \geqq \frac{r_2^3 - r_3^3}{3}.$$

Hence

$$J(\bar{C}_{12}) > \frac{r_1^3 + r_2^3}{3} = J(S_{102}).$$

We may therefore state the theorem:

THEOREM III. *If* $|\theta_2 - \theta_1| \geqq \pi/3$ *the curve* S_{102}, *formed by the straight lines* $\theta = \theta_1$, $\theta = \theta_2$, *has the least moment of inertia with respect to the point* 0 *of all curves connecting the points* 1 *and* 2.

SHEFFIELD SCIENTIFIC SCHOOL OF YALE UNIVERSITY,
NEW HAVEN, CONN.

TRIANGLES AND QUADRILATERALS INSCRIBED TO A CUBIC AND CIRCUMSCRIBED TO A CONIC

BY H. S. WHITE

PONCELET'S porism of inscribed and circumscribed polygons has a certain extension to the system of a non-singular plane cubic and a curve of the second class, resembling that recently given by Morley to point cubic and line cubic, but differing also in one feature. I shall consider only polygons of three and of four sides, first as to their possible occurrence, then by reviewing a well known construction, and lastly as to the corresponding algebraic conditions.

1. Can a triangle have vertices moving freely on a cubic while its sides continually touch a conic? If not on a general cubic, then can a special cubic admit such a system of triangles, and how strongly must it be specialized?

Call the cubic C and the conic K, and suppose that every point of C is a vertex of a triangle inscribed in C and having for sides three tangents of K. If one side of such a triangle is a, and two vertices upon that side are the points A_1 and A_2 of the cubic, let b denote a second side, and B_1, B_2, its vertices, so that A_2 and B_1 shall denote the same point. Outside of these points, a and b cut the cubic in A_3 and B_3 respectively, and these points by hypothesis must be themselves vertices of other such triangles. Through A_3, however, pass only 2 tangents of K; one is a, call the other one d. Of this second triangle, two vertices must lie on a; one is A_3, let the other be $A_2(\equiv B_1)$. Then as a and b are the two tangents to K from A_2, both must belong to this second triangle, and B_3 must be its third vertex. Similarly we should find a third triangle B_3, $B_2(\equiv C_1)$, C_3, and a fourth $C_3, C_2(\equiv A_1), A_3$. Hence *the existence of a triangle with vertices freely moving on the cubic while its sides roll upon the conic necessitates the existence of a quadrilateral with all six vertices moving freely on the cubic while its sides continually touch the conic.*

It is well known that if a complete quadrilateral has all its vertices upon the cubic, its opposite vertices are always a Steiner pair of the lowest kind, and that every cubic contains three discrete systems of such pairs. To construct the quadrilateral, two pairs may be chosen at random from any one system, then the lines joining points of one pair to points of the other constitute the quadrilateral, and its remaining vertices will fall on a third pair of the same system. Accordingly *if the desired quadrilateral can exist, its pairs of opposite vertices must be three conjugate pairs in the same one of the three systems in involution on the cubic.*

From every point of the cubic C, the pairs of points in involution on C are projected by the pairs of rays of a pencil in involution. Two rays u_1 and u_2 of each such pencil are tangent to K; replace them by their corresponding rays in the involution, v_1 and v_2. In how many pencils will it happen that v_1 coincides with u_2; i. e., that the two tangents of K from a point Z of the cubic are paired in the involution on the cubic? To answer this question, examine the whole system of lines v_1, v_2, or the v's, and determine its class.

Through Z there pass two v's conjugate to the two tangents from Z to K. But also through Z', the conjugate of Z on C, there pass two tangents of K which belong to pencils emanating from four other points; the four rays conjugate to these tangents will be additional v-lines through Z, consequently the system of v's is of class 6, and so contains $6 \times 2 = 12$ tangents of K. Some of those 12 however are improperly counted, namely those which are self-conjugate in the involutions. Self-conjugate lines (call them w-lines) fill a locus of class 3, known in certain relations as a Cayleyan envelope. Hence $3 \times 2 = 6$ of them are tangent to K, leaving but 6 tangent v-lines properly so called. As these 6 are at once u-lines and v-lines, they occur in pairs, and give only 3 points of meeting of such pairs on the cubic.

If the Steiner correspondence or involution is given, together with the cubic and conic, these 3 points are susceptible of algebraic determination, and we can infer that if any cubic contains more than 3 such points, every point upon it will satisfy the same conditions. So it is proved that

Every given cubic contains 3 points from which pass pairs of tangents to a given conic, each pair consisting of lines conjugate in one involution on the cubic. As the cubic has in general three such involutions, it will have altogether 9 such points. Further, if in one of the involutions there occur 4 such points, then every point of the cubic (except its intersections with the conic) will be the intersection of 2 conjugate tangents to the conic.

When this special case occurs, it is not difficult to show that quadrilaterals are present, of the sort described above. As one quadrilateral has six vertices, we see that the existence of one quadrilateral inscribed in the cubic and circumscribed to the conic is more than sufficient to prove the existence of an infinite system of quadrilaterals having the same property.

2. If an infinite system of Poncelet quadrilaterals is present, consider the cubic as hessian of a primitive cubic C_1; then the three diagonals of each quadrilateral will be tangents to the Cayleyan of C_1, and form a polar triangle of the conic, variable upon the Cayleyan. This gives at once a means of writing explicitly a relation between the conic and cubic, namely by using the condition that a given curve of the third class shall have tangents arranged in an infinite number of triples, each triple a polar triangle of the conic. That condition consists of the identical vanishing of a quadric covariant of the Cayleyan and the conic.*

* See *Transactions of the American Mathematical Society*, vol. 4 (1903), p. 138, (5).

As that covariant condition is incorrectly printed in its first publication,* it is desirable to give it here *in extenso*. Call the equation of the Cayleyan $u_k^3 \equiv u_{k'}^3 = 0$, and that of the conic $u_\alpha^2 = u_\beta^2 = \text{etc.} = 0$, then the condition is

$$\begin{aligned}\Gamma'(u) \equiv{}& \Delta^2(akk')^2 u_k\, u_{k'} + 3\,\Delta\,(a\beta k)(a\beta k')(\gamma\delta k)^2 u_{k'}^2 \\ &- 3(a\beta k)(a\beta k')(\gamma\delta k)^2(\epsilon\zeta k')^2.\ u_\theta^2 \\ &- \Delta\,(a\beta k)(a\beta k')(\gamma k k')^2\ u_\delta^2 \equiv 0,\end{aligned}$$

where Δ denotes the discriminant $(a\beta\gamma)^2$. This identity in variables (u) yields however but two independent conditions. These are necessary, but of course not sufficient, since we have seen that only a double infinity of conics can be found, and accordingly *three* conditions are to be imposed. Such a set is found in the following manner.

It is well known that in a doubly infinite linear system (or web) of line-conics, four common tangents to any two of the conics will intersect in six points of a fixed point-cubic. The pairs of opposite points in any such quadrilateral are Steiner pairs of the lowest kind upon the point-cubic. That cubic is Cayleyan to a certain line-cubic $\Pi(u) = 0$, whose system of quadric polars consists of the line-conics in the web first mentioned; and if the point-cubic be given, the line-cubic $\Pi(u) = 0$ is one of three determinate curves, so that we find three doubly infinite webs of conics standing in the required relation. Probably the simplest statement is obtained by assuming as known a point-cubic, $f = 0$, of which the given cubic, $\Delta^3(x) = 0$, is hessian. The form $f(x)$ determines uniquely a doubly apolar form $\Pi(u)$, whose first polars are then the conics about which quadrilaterals can be circumscribed so as to move freely while inscribed to the hessian. Analytically the conditions can be written, first in terms of a single one of the three pro-hessians, $f(x) = 0 = f_x^3$, of the given cubic $\Delta_x^3 = 0$.

$$\begin{aligned}&1)\quad (\Delta\Delta' u)^2 \Delta_\phi \Delta'_\phi - \tfrac{1}{6}\, S(ff'u)^2 f_\phi f'_\phi \equiv \quad \mu \cdot u_\phi^2, \\ &2)\quad (\Delta\Delta' u)^2 \Delta_\psi \Delta'_\psi - \tfrac{1}{6}\, S(ff'u)^2 f_\psi f'_\psi \equiv -\mu \cdot u_\psi^2, \\ &3)\quad f_x^2 f_y \equiv 0.\end{aligned}$$

In these defining equations, (1) and (2) are identities in variables (u), S is the invariant of degree 4 in the coefficients of f_x^3, μ and $-\mu$ are constants whose

* The explicit reckonings following p. 138, loc. cit., are correct, the only errors being in formulæ (4) and (5).

value depends on the cubic f_x^3 alone,* and $u_\phi^2 = 0$, $u_\psi^2 = 0$, $u_\chi^2 = 0$ denote the conics of the three systems respectively; the identity (3) is in variables (y). These equations have the merit of being linear in the coefficients of the conic. All three systems may be defined by equations like (3) if we assume as known the three pro-hessians $f_x^3 = 0$, $g_x^3 = 0$, and $h_x^3 = 0$ of the given cubic $\Delta_x^3 = 0$. The conditions are then:

$$f_\chi^2 f_y \equiv 0,\ g_\phi^2 g_y \equiv 0, \text{ and } h_\psi^2 h_y \equiv 0.$$

These are well known systems, but this their peculiar property seems not to have been exhibited as an extension of the theorems connected with Poncelet's polygons.† Schroeter's celebrated linear construction of conjugate point pairs on the cubic from three given pairs requires the drawing of exactly the lines which have been described as tangents to conics, four at a time giving rise to one new pair of vertices. Calling these a Schroeter quadrilateral, we may formulate what has been proved:

A cubic curve being determined by Schroeter's construction from three pairs of points, the eight sides of any two Schroeter quadrilaterals are tangent to some conic, which envelopes a simply infinite system of such quadrilaterals; and all such conics form a doubly infinite linear system of line-conics, the first polars of a line-cubic which has Schroeter's cubic for its Cayleyan.

Given one Schroeter quadrilateral and two conjugate points not upon its sides, another such quadrilateral can be constructed by solving a single problem of the second order, or a quadratic equation.

VASSAR COLLEGE, POUGHKEEPSIE, N. Y.
OCTOBER, 1905.

---

* See Conics and Cubics connected with a plane cubic etc., *Transactions of the Amer. Math. Society*, vol. 1 (1900), p. 8.

† See however Emil Weyr, Uber Involutionen höherer Grade, *Crelle's Journal*, vol. 72 (1870), pp. 285–292, and S. Kantor, Verallgemeinerungen eines Poncelet'schen Satzes, *Crelle's Journal*, vol. 86 (1879), p. 271.

ON A FUNCTION WHICH OCCURS IN THE LAW OF THE MEAN*

By E. R. Hedrick

1. Introduction.

The formula

$$f(x+h) - f(x) = h \cdot f'(x+\theta h) = h \cdot f'(x+\xi), \qquad 0 < \theta < 1,$$

is known to every student of mathematics under the title "the law of the mean" or a synonym. It is usually proved that the formula is correct if $f(x)$ is defined in an interval $a \leqq x \leqq b$ and if the derivative $f'(x)$ exists in the continuum $a < x < b$.†

The quantity ξ which occurs in the formula is evidently a function of the two independent variables x and h and is defined for all values of x and h for which x and $x + h$ both lie in the interval in which $f'(x)$ exists. The purpose of the present paper is to discuss the properties of this quantity ξ, and we shall write

$$\xi = \xi\,(x, h)$$

whenever it is desired to emphasize its functional character.

In studying the function $\xi(x, h)$ it occurs to one immediately that if $\xi(x, h)$ is continuous, the derivative $f'(x)$ is surely continuous (see p. 183). A slight inspection would tend to convince one that the converse is true—at least in a limited sense—i. e., that if $f'(x)$ is continuous, $\xi(x, h)$ is a continuous function of h when x is constant. This conclusion is, however, erroneous. It will be shown that $\xi(x, h)$ may be utterly discontinuous, in the sense that ξ fails to take on values which lie between values which it does take on, even when $f'(x)$ exists and is continuous everywhere (see p. 190).

The fallacies which depend upon the utter discontinuity of ξ are known, but not well-known. While pointing out their dangers, it may be well to remark that certain proofs may be conducted without error along the same general lines as these fallacious arguments, if the function ξ be fully understood.

Incidentally it will become evident that the law of the mean may hold in its present form in case the derivative $f'(x)$ does not even exist at certain points, even when we permit x and $x + h$ to assume any values whatever; and generalized statements of the law hold even in more tenuous cases.

---

* This paper is based partially upon a paper entitled "On the Law of the Mean," read by the writer at the summer meeting, 1903, of the American Mathematical Society, and partially on another paper entitled "The function $\xi(h)$ in the Law of the Mean," read by the writer at the April meeting, 1906, of the Chicago Section of the Society.

† A proof of this fact will incidentally occur as a corollary to a theorem of this paper.

Finally, certain unusual examples in the theory of maxima and minima which are suggested by the paper will be presented and commented upon for their useful warning against too great assumption in that theory.

2. Limits over an assemblage. *A function $f(x)$ of a real variable x is said to approach a limit K as x approaches a particular value a if, corresponding to any pre-assigned positive number ${}^1\epsilon$, another positive number ${}^2\delta$ exists, such that*

$$(1) \qquad |f(x) - K| < {}^1\epsilon \quad \text{whenever} \quad 0 < |x - a| < {}^2\delta$$

*where x may be any number whatever which satisfies the last inequality.** I shall say for distinctness and emphasis that $f(x)$ approaches K *with respect to the continuum* or *in the ordinary sense* whenever these conditions are fulfilled. The left-handed superscripts indicate the order of choice of ϵ and δ.

It may happen that the state of affairs described by (1) holds only in case x is allowed to assume the values which belong to a certain assemblage (E). In this case I shall say that $f(x)$ approaches K *with respect to the assemblage* (E) or simply *over* (E), and I shall write

$$\lim_{x=a \ (E)} f(x) = K.$$

It is evidently trivial to consider a case in which a is not a limiting point of (E), since any function evidently approaches any desired constant for such an assemblage. We shall therefore assume that $x = a$ is a limiting point of (E) unless the contrary is stated.

In particular, if (E) consists of all points to the left (right) of a, and if the preceding inequalities are satisfied when x takes on any value in (E), *then $f(x)$ will be said to approach K from the left (right).*

If $f(a) = K$ and if $f(x)$ approaches K with respect to an assemblage (E) as x approaches a, we shall naturally say that $f(x)$ *is continuous at a with respect to* (E) or simply *over* (E).

It is now very easy to write down the following theorems, at least some of which are important:—

* It should be noted, especially for what follows, that this definition absolutely eliminates any consideration of *time*; nothing whatever is said concerning the order in which the values of x are taken on, and all notion of succession is eliminated from the vulgar notion "approach."

I. *Let (E) be an assemblage with a single limiting point a. Then (E) is countable and its members may be written in a sequence*

$$(E): \quad x_1, x_2, x_3, \cdots, x_n, \cdots;$$

if $f(x)$ approaches K over (E) we shall have $\lim_{i=\infty} f(x_i) = K$. For ${}^2N$ can be chosen so large that $|x_n - a| < {}^1\epsilon$ whenever $n \geqq {}^2N$.

II. *If $f(x)$ approaches a limit K with respect to any assemblage (E), it approaches K with respect to any sub-assemblage of (E).*

III. *If $f(x)$ approaches K over (E), the assemblage (E) may be increased by adjoining the points of any assemblage for which $x = a$ is not a limiting point.*

IV. *More generally, if $f(x)$ approaches K over each of a finite number of assemblages $(E)_1, (E)_2, \cdots, (E)_n$, the limit is the same over the assemblage (E) formed by uniting all the $(E)_i$.* For, given ϵ, a δ can be found for each $(E)_i$; the least of these δ's may be used for (E).

V. *In particular, if the finite set of assemblages $(E)_i, (i = 1, 2, \cdots n)$, exhaust all possible numbers in the neighborhood of $x = a$, (E) is the continuum, and $f(x)$ is continuous in the ordinary sense.*

VI. *On the other hand, if $f(x)$ approaches K with respect to any assemblage (E), any number whatever $b \neq a$ is a member of some assemblage (F) for which $f(x)$ approaches K.* For at least one such (F) is formed by adjoining b to (E).

VII. *It follows that even though $f(x)$ approaches K for each of a set of assemblages (E) which exhaust all possible numbers, $f(x)$ does* NOT *necessarily approach K in the ordinary sense.* This follows directly from VI and from the knowledge of a single example of a function which approaches no limit in the ordinary sense, but which does approach a limit over some one assemblage. Such an example is $y = f(x) = \sin(1/x)$; this function approaches no limit as x approaches zero, in the ordinary sense, but approaches the limit zero over the assemblage $(E) = 1/n\pi$ $(n = 1, 2, \cdots)$. Hence, by VI, $f(x)$ approaches zero over each of a set of assemblages which exhaust all possible numbers. Even if the set of (E)'s for which $f(x)$ approaches K and which exhaust all possible numbers is known to be a countable (or enumerable) infinite set, we cannot conclude that $f(x)$ approaches K in the ordinary sense. [The example $f(x) = 0$ when x is irrational, $f(x) = i \cdot x$ when $x = \frac{p}{q^i}$, where p is prime to q^i and q is not a perfect power, p, q, i being all positive integers,

is an example of this kind. For $\lim f(p/q^i) = 0$ for any fixed value of i, and also $\lim f(x) = 0$ if x is irrational, but $f(x)$ does not approach zero in the ordinary sense, as is seen by actually drawing the figure.*] *If the approach is uniform, however, the conclusion that $f(x)$ approaches K is correct.*

VIII. There is another theorem which is applicable in cases not covered by V. For *if we know that $f(x)$ approaches K for every possible assemblage, or even if we know merely that $f(x)$ approaches K for every possible sequence, we may conclude that $f(x)$ approaches K in the ordinary sense.*† For, if $f(x)$ does not approach K, there exists a positive number ϵ for which relations (1) are false for every δ. Choosing $\delta_i < 1/2^i$, we shall then have $|f(x_i) - K| > \epsilon$ for some one $x_i < \delta_i$. Then $f(x)$ surely does not approach K over the sequence of the x_i's.

These results evidently apply also in testing a function for *continuity over an assemblage* (E) at a point $x = a$, provided that $f(a) = K$.

Again, we may substitute the notion of *oscillation* for that of approaching a limit. If we consider all the points of an assemblage (E) which lie in a variable interval about $x = a$, it is evident that there is one and only one lower limit of the upper limits of $f(x)$ on (E) in all such intervals. Let us call this L. Likewise there is only one upper limit of the lower limits of $f(x)$ on (E) in all such intervals, which we call l. Then *the oscillation at $x = a$ with respect to (E) is* $\omega_{(E)} = L - l$, and it is evidently necessary and sufficient that $\omega_{(E)} = 0$ in order that $f(x)$ be continuous at $x = a$ with respect to (E). This notion may be generalized essentially;‡ and theorems analogous to those above may be written down.

3. The function $\xi(x, h)$. If $f(x)$ be defined in the interval $a \leqq x \leqq b$, the quotient

$$Q(x, h) = \frac{f(x + h) - f(x)}{h}$$

is a function of x and h defined for values of x and h for which x and $x + h$ both lie in the interval $a \leqq x \leqq b$. If this function $Q(x, h)$, for a constant value of x, approaches a limit K(as h approaches zero) over any assemblage

* See also *Bull. Amer. Math. Soc.*, vol. 11, footnote, p. 321, March, 1905.

† A generalization of this theorem is obvious: If $f(x)$ approaches K for every possible sequence which can be formed from the members of any assemblage (E), then $f(x)$ approaches K over (E).

‡ I shall do this in another paper.

(E) for which $h = 0$ is a limiting point, *K is said to be the derivative of $f(x)$ with respect to (E)*, or simply *over (E)*. If (E) is the continuum, the derivative $f(x)$ exists *over the continuum*, or simply *in the ordinary sense*. Otherwise we shall write

$$K = D_{(E)} f(x).$$

Let us consider the auxiliary function

$$\phi(x) = f(x) - f(a) - \frac{f(b) - f(a)}{b - a}(x - a).$$

Let us suppose that $\phi(x)$ is at a maximum (or at a minimum) at some point $x = c$, $a < c < b$. This will certainly be the case, for example, if $f(x)$ is continuous, since $\phi(b) = \phi(a) = 0$. Then if $D_{(E)}\phi(c)$ exists, where (E) denotes any essentially two-sided assemblage,* we must have

$$D_{(E)}\phi(c) = D_{(E)} f(x) - \frac{f(b) - f(a)}{b - a} = 0.$$

IX. *Hence if $\phi(x)$ is at a maximum (or at a minimum) at $x = c$, $a < c < b$ (in particular, if $f(x)$ is continuous) and if $D_{(E)} f(c)$ exists for a two-sided assemblage (E), we shall have*

$$f(b) - f(a) = (b - a) D_{(E)} f(c), \qquad a < c < b.$$

This is the law of the mean. The requirement that $f(x)$ be continuous in the interval $a \leqq x \leqq b$ and that the ordinary derivative $f'(x)$ exist for $a < x < b$ is evidently *sufficient*, but is *in no sense necessary* for this result.

Replacing a by x and $b - a$ by h, the formula may be written in the form

$$Q(x, h) = \frac{f(x + h) - f(x)}{h} = f'(x + \xi), \qquad 0 < \xi/h < 1,$$

at least if $f'(x)$ exists throughout an interval in which x and $x + h$ lie. That this formula holds in a very general manner in the precise form just given, even when $f(x)$ is discontinuous between x and $x + h$, is evident from such an example as $y = f(x) = \sin(1/x)$, which is essentially discontinuous at $x = 0$. If we add $f(0) = 0$, the function $f(x)$ is defined for every value of x, and it is easy to see that the preceding formula holds when x and $x + h$ assume absolutely any values whatever.

* That is, an assemblage in which there are points on both sides of $x = a$ in any neighborhood of $x = a$.

We shall now assume, however, that $f'(x)$ exists in the ordinary sense. The number ξ depends upon the choice of both x and h, and may have several values, or even an infinite number, for one pair of values of x and h. If x is fixed, ξ has at least one value for every value of h and that value is numerically less than h; i. e., in a plane whose coordinates are ξ and h, ξ always lies between the h-axis and the line $\xi = h$.

There is a common fallacy which we may revise into an accurate proof. Since $f'(x)$ exists, we shall have, at any point $x = k$,

$$\lim_{h=0} Q(k, h) = f'(k)$$

and since

$$Q(k, h) = f'(k + \xi), \qquad 0 < \xi/h < 1$$

we shall also have

$$\lim_{h=0} f'(x + \xi) = f'(k).$$

Since ξ is numerically less than h, $\xi(k, h)$ approaches zero with h: and we might be led to infer that

$$\lim_{\xi=0} f'(k + \xi) = f'(k),$$

i. e., that $f'(x)$ is continuous at $x = k$. *This is false*, however; and the fallacious nature of the supposed conclusion is exposed by a single well-known example in which $f'(x)$ exists and is discontinuous:

$$y = f(x) = x^2 \sin(1/x^2) \qquad \text{if } x \neq 0,$$
$$y = f(x) = 0 \qquad \text{if } x = 0.$$

The function $f(x)$ thus defined is evidently continuous, and its derivative is

$$y' = f'(x) = 2x \sin(1/x^2) - (2/x)\cos(1/x^2) \qquad \text{if } x \neq 0,$$
$$y' = f'(x) = \lim_{h=0} h \sin(1/h^2) = 0 \qquad \text{if } x = 0.$$

These forms show that $f'(x)$ exists for every value of x, and that $f'(x)$ is discontinuous at $x = 0$, for $f'(x)$ exceeds any fixed number in any interval about $x = 0$. This example illustrates also the correct conclusions which follow.

In fact the function $\xi(0, h)$ does not assume all values near zero for values of h which lie in *any* limited interval whatever. For $|f'(x)|$ exceeds any fixed constant [2]C inside of any interval $(-{}^1\epsilon, 0)$ and also inside $({}^1\epsilon, 0)$. Such is the real nature of the fallacy in any case: for *if* $\xi(k, h)$ *is a continuous function of* h, or in fact even *if having assigned* [1]ρ *we can find* [2]λ *such that* $|\xi|$

takes on all values less than λ *for some values of* $|h|$ *less than* ρ, *the preceeding argument really shows that* $f'(x)$ is *continuous at* $x = k$. For, having chosen $\epsilon_1, \epsilon_2, \cdots, \epsilon_n, \cdots$ $(\lim_{i=\infty} \epsilon_i = 0)$ we can choose $\rho_1, \rho_2, \cdots, \rho_n, \cdots$ $(\lim_{i=\infty} \rho_i = 0)$ in such a way that

$$|Q(k, h) - f'(k)| < \epsilon_i, \text{ whenever } |h| < \rho_i.$$

If, corresponding to each ρ_i a quantity λ_i can be found such that $|\xi|$ takes on every positive value less than λ_i for some value of $|h|$ less than ρ_i we shall have

$$|Q(k, h) - f'(k)| \equiv |f'(k + \xi) - f'(k)| < \epsilon_i \text{ whenever } |\xi| < \lambda_i,$$

which expresses that $f'(x)$ is continuous at $x = k$.

X. *A sufficient condition that* $f'(x)$ *should be continuous is that corresponding to any positive number* $^1\rho$ *another positive number* $^2\lambda$ *exists, such that* $|\xi|$ *takes on all values less than* λ *for some value of* $|h|$ *less than* ρ, *for both positive and negative values of* ξ.

Geometrically, a sufficient condition that a variable tangent should approach a fixed tangent is that in any interval about the fixed point of tangency a subinterval exists such that any tangent in the subinterval is parallel to some secant in the original interval through the fixed point.

Whether this condition be satisfied or not, we can always find *an assemblage* (E) *with respect to which* $f'(x)$ is continuous at $x = k$. For, having chosen h_1, there exists at least one corresponding ξ different from zero, say ξ_1. Setting $h_2 = \xi_1$, we can find a corresponding $\xi_2 \neq 0$. Proceeding in the same manner we find ξ_i, $i = 1, 2, \cdots$, $|\xi_{i+1}| < |\xi_i|$, where $\lim_{i=\infty} \xi_i = 0$. Since $Q(k, h)$ approaches $f'(k)$ as h approaches zero, $\lim_{i=\infty} Q(k, h_i) = f'(k)$, and therefore

$$\lim_{i=\infty} f'(k + \xi_i) = \lim_{i=\infty} Q(k, h_i) = f'(k),$$

which shows that $f'(x)$ approaches $f'(k)$ over $(E) = k + \xi_1, k + \xi_2, \cdots$ as x approaches k. Hence we may say

XI. *A derivative which exists at every point in an interval is continuous at each point of the interval with respect to at least one essentially two-sided assemblage.*

XII. *It follows that if the derivative* $f'(x)$, *which is supposed to exist at every point, approaches any limit whatever in the ordinary sense, as* x *approaches* k, *that limit is* $f'(k)$.

XIII. *It is evident from* VI *that if* $f'(x)$ *exists near* $x = k$ *the assemblages with respect to which* $f'(x)$ *approaches* $f'(k)$ *as* x *approaches* k *exhaust all possible numbers.* But there are still other assemblages for which $f'(x)$ does not approach $f'(k)$ unless $f'(x)$ is continuous at $x = k$. (See VII).

It is easy to generalize these statements somewhat. For the derivative $f'(x)$, which is supposed to exist, surely takes on any value C $(A < C < B)$ at some point $x = t$ $(a < t < b)$ if it takes on the values A and B at $x = a$ and $x = b$, respectively. For, $Q(x, h)$ being continuous in both x and h, $h_0 > 0$ can surely be choosen so small that $Q(a, h_0) < C$ and $Q(b - h_0, h_0) > C$. Hence $Q(c, h_0) = C$, since Q is continuous in x. Then, by the law of the mean $Q(c, h_0) = f'(c + \xi) = C$, where $s = c + \xi$ lies between a and b since $h_0 > \xi > 0$ and $a < c < b - h_0$. This fact is well known, i. e., *in passing from one value to another, any derivative* $f'(x)$ *takes on every intermediate value provided it exists at every point.*

XIV. *It follows that if* $f'(x)$ *approaches two different limits* m *and* n *(as* x *approaches* k*) with respect to two different assemblages* (M) *and* (N)*, then* $f'(x)$ *approaches any intermediate value* p*,* $(m < p < n)$*, with respect to some third assemblage* (P). For $f'(x)$ assumes values which differ from m (and also values which differ from n) by less than any preassigned number ϵ near $x = k$; hence $f'(x)$ actually assumes the value p at an infinite number of points on both sides of $x = k$, which proves the theorem. In particular, *if* $f'(x)$ *approaches any other value* $V \neq f'(k)$ *over any assemblage, then* $f'(x)$ *also approaches any number between* V *and* $f'(k)$ *over some assemblage.* The property mentioned in the theorem is another property of continuous functions which is shared by any existing derivative, for if $\phi(x)$ is continuous except at $x = k$, $\phi(x)$ surely has the property in question. We shall resume this discussion presently.

Again, using the previous notation (p. 183) let ξ_i and ξ_{i+1} be the values of ξ which correspond to h_i and h_{i+1}, respectively, and let $h_i = \xi_{i+1}$. Then we shall have $Q(k, h_i) = f'(k + \xi_i) = L_i$, say, for each value of i. But since $Q(k, h)$ is continuous in h, it assumes every value between L_i and L_{i+1} for some value of h between h_i and h_{i+1}. Hence the assemblage of values of h between h_i and h_{i+1} for which $Q(k, h)$ lies between L_i and L_{i+1} *has the power of the continuum.* Since $f'(x)$ also assumes all intermediate values in passing from one value to another, $f'(k + \xi)$ also assumes every values between L_i and L_{i+1} for some value of h between h_i and h_{i+1}. Hence for the values of h found above we shall have $Q(k, h) = f'(k + \xi)$ where h lies between h_i and h_{i+1} and

where ξ lies between ξ_i and ξ_{i+1}. Thus the assemblage of values of ξ which lie between ξ_i and ξ_{i+1} and which correspond to values of h between h_i and h_{i+1} is an assemblage *which has the power of the continuum*. Let us then consider the sequence $\xi_1, \xi_2, \cdots, \xi_n, \cdots$ and also the interpolated values of ξ just found. The total assemblage (T) formed by uniting all these has the power of the continuum in any interval about $\xi = 0$. On the other hand it is evident that $f'(k + t)$ approaches $f'(k)$ with respect to this assemblage (T). It follows that

XV. *If $f'(x)$ exists in any interval, it is continuous at every point of that interval with respect to a certain assemblage (T) which has the power of the continuum on each side of the point in any interval whatever about the point.*

It would be fallacious to conclude that $f'(x)$ is continuous at any point with respect to the assemblage of values which $k + \xi$ may assume, and this conclusion might lead to as grave error as the previous fallacy. For example, in the previous exercise $y = f(x) = x^2 \sin(1/x^2)$, ξ may actually take on any value whatever for some value of h. Hence the conclusion that $f'(x)$ is continuous for the assemblage of all possible $k + \xi$ would lead to the erroneous conclusion that $f'(x)$ is continuous in the ordinary sense.

4. Derivatives as limits of continuous functions. The function $Q(x, h)$ approaches $f'(x)$, if $f'(x)$ exists, for any sequence of values of h whose limit is zero. Selecting such a sequence:

$$1) \qquad h_1, h_2, \cdots, h_n, \cdots$$

where the h's do not depend on x,* the corresponding values of Q, viz:

$$2) \qquad Q(x, h_1),\ Q(x, h_2), \cdots, Q(x, h_n), \cdots$$

form a sequence of continuous functions of x, and the series

$$3) \quad Q(x, h_1) + [Q(x, h_2) - Q(x, h_1)] + [Q(x, h_3) - Q(x, h_2)] + \cdots + [Q(x, h_{i+1}) - Q(x, h_i)] + \cdots$$

is a series of continuous functions which surely converges to the sum $f'(x)$ for any value of x for which $f'(x)$ exists. The results concerning such series are therefore immediately applicable. If the series (3) converges uniformly, for example, the derivative $f'(x)$ is continuous, for the sum of a uniformly con-

* It is easy to see that most of what follows is independent of this assumption.

vergent series of continuous functions is continuous. Such is the case, in particular, if

$$|f'(x) - Q(x, h_i)| < {}^1K \qquad \text{whenever } i > {}^2N;$$

or again if

$$|Q(x, h_n) - Q(x, h_p)| < {}^1K \qquad \text{whenever } a > n > {}^2N,$$

where N is independent of x. That is to say, if $Q(x, h)$ approaches its limit $f'(x)$ *uniformly* for any constant sequence 1) and for all values of x, then $f'(x)$ is continuous. Any example of a derivative which exists and is discontinuous (e. g., the example given above) is also an example of a series of continuous functions whose limit exists and is discontinuous, i. e., an example of non-uniform approach; and that for any sequence of h's of the type 1).

On the other hand, suppose that $f'(x)$ is continuous. Then since any continuous function is uniformly continuous, $f'(x + t)$ approaches $f'(x)$ uniformly. Therefore

$$|f'(x) - Q(x, h_i)| < {}^1K \text{ whenever } i \geqq {}^2N, \text{ independent of } x,$$

for $Q(x, h_i) = f'(x + \xi_i)$ and $f'(x + \xi_i)$ surely approaches $f'(x)$ uniformly. It follows that if $f'(x)$ is continuous the sequence 2) is uniformly convergent; hence the series 3) [and also the sequence 2)] has the following unusual property:

XVI. *The sum $f'(x)$ of the series 3) of continuous functions is continuous in an interval when and only when the series 3) converges uniformly in that interval.*

It is shown by Baire* that the limit of any sequence of continuous functions must be continuous at least once (and therefore an infinite number of times) in any interval in which the limit exists, and the obvious conclusion is drawn that any derivative which exists throughout an interval is continuous at at least one point in every subinterval.

If the derivative $f'(x)$ is discontinuous at any poiut $x = a$, it must be possible to assign a positive number ϵ such that $|f'(x) - f'(a)| > \epsilon$ for some x in any neighborhood of $x = a$. But since $f'(x)$ takes on all intermediate values between any two which it does take on, it follows that at any point of discontinuity $x = a$ a positive number ϵ depending upon a can be assigned, such that $|f'(x)|$ takes on every value between $|f'(a)|$ and $|f'(a)| + \epsilon$ in any neighborhood of $x = a$.

* Baire, Thesis: *Sur les fonctions discontinues*, p. 30 and p. 108.

XVII. *The range of values all of which a derivative actually takes on in any neighborhood of any point $x = k$ is zero only when the derivative is continuous at $x = k$.*

Let us inquire whether a derivative may be discontinuous at points which are everywhere dense in an interval throughout which it exists. Any discontinuous derivative (e. g., the example given above) is an example of the remarkable class of functions studied by Darboux, which are not continuous, but which do assume all intermediate values. It is even possible to construct a function which is discontinuous at every point of an interval but which assumes all intermediate values in passing between any two which it does assume.* Such a function is not the derivative of any function whatever, on account of Baire's theorem.

On the other hand, it is easy to write down a function which is continuous at points everywhere dense and also discontinuous at points everywhere dense; e. g., the well known function

$$F(x) = 0 \quad \text{if } x \text{ is irrational}; \quad F(x) = 1/q \quad \text{if } 0 < x = p/q < 1$$

is defined whenever $0 < x < 1$, is continuous at every irrational point and discontinuous at every rational point in that interval. This function is not the derivative of any function whatever, since it does not assume all intermediate values in the necessary manner.

The question remains then whether it is possible to construct a function which has all these peculiar properties, i. e., a function such that the range of values all of which it actually takes on in the neighhorhood of a point is zero at some point and different from zero at some other point in every interval. At first sight this seems impossible, but it becomes evident that such functions actually exist if we first write the theorem:

---

* See Lebesgue, *Leçons sur l'integration* (1905), p. 90. Lebesgue defines $\phi(x)$ for $0 < x < 1$ as follows: having written x as an ordinary decimal $x = \frac{a_1}{10} + \frac{a_2}{10^2} + \frac{a_3}{10^3} + \ldots$ he considers the sequence $a_1, a_3, a_5, a_7, \ldots$; if this sequence is *not* periodic he defines $\phi(x) = 0$; if the same sequence is periodic, if a_{2n-1} is the first member of a period, he defines

$$\phi(x) = \frac{a_{2n}}{10} + \frac{a_{2n+2}}{10^2} + \frac{a_{2n+4}}{10^3} + \ldots$$

Then $\phi(x)$ takes on every value between 0 and 1 in every subinterval of the interval $0 \leqq x \leqq 1$; but $\phi(x)$ is discontinuous at every point.

XVIII. *Either* (*a*) *the derivative of any differentiable function is continuous throughout some subinterval of any interval in which it exists;* or else (*b*) *the derivative has the unusual property just mentioned.*

The discovery of a derivative which exists at every point and which is discontinuous at points everywhere dense in any interval would therefore lead to a remarkable example of a function. It is obvious that such a function results if we apply Hankel's principle of condensation of singularities to the preceding example $x^2 \sin(1/x^2)$, for we can write down at once a uniformly convergent series which has a derivative everywhere, which itself is discontinuous at points everywhere dense.* *Thus there actually exists functions* (*e. g.*,

---

* Such examples are known. The intention of this remark is to call attention to the consequence mentioned above.

Lebesgue indicates (l. c., p. 94), but does not actually write out, the following example:

Let $$\left.\begin{aligned} \phi(x, a) &= (x-a)^2 \sin \frac{1}{x-a}, & x \neq a \\ &= 0, & x = a \end{aligned}\right\} \quad \begin{matrix} 0 \leqq a \leqq 1 \\ 0 \leqq x \leqq 1 \end{matrix}$$

and let

$\phi_i(x) = \phi(x, a_i)$ where $a_1, a_2, \ldots, a_n, \ldots$ are the elements of any countable dense assemblage (E) (e. g., all rational numbers) arranged in a sequence.

Then the series

$$\phi(x) = \phi_1(x) + \frac{1}{2!}\phi_2(x) + \ldots + \frac{1}{n!}\phi_n(x) + \ldots, \quad 0 \leqq x \leqq 1 \tag{1}$$

is an absolutely and uniformly convergent series, $0 \leqq x \leqq 1$.

Let $$\left.\begin{aligned} \psi(x, a) = \frac{\partial \phi(x, a)}{\partial x} &= 2(x-a) \sin \frac{1}{x-a} - \cos \frac{1}{x-a}, & x \neq a \\ &= 0, & x = a \end{aligned}\right\} \quad \begin{matrix} 0 \leqq a \leqq 1 \\ 0 \leqq x \leqq 1 \end{matrix}$$

and let $\psi_i(x) = \psi(x, a_i)$. Then the series of discontinuous functions:

$$\psi(x) = \psi_1(x) + \frac{1}{2!}\psi_2(x) + \ldots + \frac{1}{n!}\psi_n(x) + \ldots \tag{2}$$

converges absolutely and uniformly, $0 \leqq x \leqq 1$, and the individual terms, $\psi_i(x)$, are the derivatives of the individual terms of (1). *It follows that $\psi(x)$ is the derivative of $\phi(x)$, and that $\psi(x)$ is discontinuous at each point of* (E).

In order to prove this we need to show first that (2) for every value of x is a uniformly convergent series of continuous functions plus at most a single discontinuous function; hence $\psi(x)$ is continuous except at points of (E) and is discontinuous at any point a_i of (E) with the same discontinuity as $\frac{1}{i!}\psi(x, a_i)$.

We must also show that $d\phi(x)/dx = \psi(x)$. *This follows from the fact that* (2) *is a uniformly convergent series of derivatives of the terms of* (1) although $\int \psi(x)dx$ does not exist in Riemann's sense.

$\psi(x)$ in footnote) which are (1) discontinuous at some point in every interval, (2) continuous at some (other) point in every interval, and which (3) actually take on every intermediate value between any two values which they do assume.

5. The Derivative Continuous. Let us now consider a function $f(x)$ whose derivative exists and is continuous throughout some interval $a \leqq x \leqq b$. We know that a sufficient condition for the continuity of $f'(x)$ is that $\xi(x, h)$ be a continuous function of h when x is constant, or even that $|\xi|$ assume all values less than $^2\lambda$ for some value of $|h|$ less than $^1\rho$. Conversely, *can we infer that if $f'(x)$ is continuous, it will follow that $\xi(x, h)$ is a continuous function of h when x is constant? Or even that $|\xi|$ assumes all values less than $^2\lambda$ for some value of $|h|$ less than $^1\rho$?* Such a conclusion seems to be obvious at first sight, but it is untrue.

Let us consider, for example, the function

$$y = f(x) = x^3\left(1 + \sin\frac{1}{x}\right) \qquad \text{if } x \neq 0,$$
$$y = f(x) = 0 \qquad \text{if } x = 0.$$

The derivative is

$$y' = f'(x) = 3x^2\left(1 + \sin\frac{1}{x}\right) - x\cos\frac{1}{x} \qquad \text{if } x \neq 0,$$
$$y' = f'(x) = 0 \qquad \text{if } x = 0,$$

which is continuous for all values of x. Nevertheless there are values of $\xi(0, h)$ which are not taken on for any value of x whatever and which lie as close to zero as we please. For it is obvious that

$$f'(x) = 0 \quad \text{when } x = \frac{2}{(4n+1)\pi}$$

and that

$$f(x) = 0 \quad \text{when } x = \frac{2}{(4n+1)\pi};\ f(x) > 0 \quad \text{when } x \neq \frac{2}{(4n+1)\pi}.$$

Now $f'(x) \not\equiv 0$, since $f'(x) \not\equiv$ const. Moreover $f'(x) \geqq 0$ is surely false, since $f(0)$ is not a minimum to the right. Hence $f'(x) < 0$ for some value of x, say x_n, which lies between $\frac{2}{(4n+1)\pi}$ and $\frac{2}{(4n+5)\pi}$. It follows that ξ never takes on the value x_n, since in the formula

$$f(h) - f(0) = f'(\xi), \qquad 0 < \xi < h$$

the right-hand side is never negative. We may therefore conclude that

XIX. *It is entirely possible that $f'(x)$ exists and is continuous throughout an interval about a point $x = k$, and yet that the assemblage of values which the function $\xi(k, h)$ never assumes has the power of the continuum.* For since $f'(x)$ is continuous, it assumes negative values (say) for all values of x in the neighborhood of any point where it is negative.

6. Maxima and Minima. Some of the remarks made above and several of the examples are suggestive in the theory of ordinary maxima and minima. It follows directly from the law of the mean that *if*

$f'(a) = 0$, *and* $f'(a + h) < 0$, *and* $f'(x - h) > 0$, *whenever* $0 < h < \delta$ *then* $f(x)$ *is at a proper maximum* * *for* $x = a$. It seems at first sight obvious that the converse would also be true and such a statement can be found without extensive search. Let us examine a few problems.

In the example

$$y = f(x) = x^2\left(1 + \sin\frac{1}{x^2}\right) + e^{-\frac{1}{x^2}} \qquad \text{if } x \neq 0,$$

$$y = f(x) = 0 \qquad \text{if } x = 0,$$

which is a slight modification of that on p. 182, the derivative is

$$y' = f'(x) = 2x\sin\frac{1}{x^2} - \frac{2}{x}\cos\frac{1}{x^2} + 2x + \frac{2}{x^3}e^{-\frac{1}{x^2}} \qquad \text{if } x \neq 0,$$

$$y' = f'(x) = 0 \qquad \text{if } x = 0.$$

Since the term $(2/x)\cos(1/x^2)$ determines the sign of $f'(x)$ for values of x for which $|\cos(1/x^2)|$ is sufficiently large near $x = 0$, it is evident that $f'(x)$ takes on both positive and negative values on each side of $x = 0$ inside of any interval about $x = 0$. It follows that the preceding method leads to no result. Moreover $f''(x)$ does not even exist for $x = 0$. Hence the only usual method for determining whether $x = 0$ gives a minimum is by examining the function directly. It is evident, however, that $f(x)$ is at a proper minimum at $x = 0$.

An objection might be made that $f'(x)$ is discontinuous at $x = 0$ in the preceding example. To meet such reasonable objection, we may cite the example

$$y = f(x) = x^4\left(1 + \sin\frac{1}{x}\right) + e^{-\frac{1}{x^2}} \qquad \text{if } x \neq 0,$$

$$y = f(x) = 0 \qquad \text{if } x = 0,$$

* That is $f(a) < f(a \pm h)$, $0 < h < \delta$. If merely $f(a) \leqq f(a \pm h)$ the minimum is said to be *improper*.

which is a slight modification of the example of p. 189. Here the derivative is

$$y' = f'(x) = 4x^3\left(1 + \sin\frac{1}{x}\right) - x^2\cos\frac{1}{x} + \frac{2}{x^3}e^{-\frac{1}{x^2}} \qquad \text{if } x \neq 0,$$

$$y' = f'(x) = 0 \qquad \text{if } x = 0,$$

which is continuous for all values of x. But for values of x which satisfy

$$0 < \bar{x} = \frac{1}{2n\pi}$$

we find

$$\cos(1/\bar{x}) = 1, \quad \sin(1/\bar{x}) = 0, \quad f'(\bar{x}) = \bar{x}^2\left(4\bar{x} - 1 + \frac{2}{\bar{x}^5}e^{-\frac{1}{\bar{x}^2}}\right).$$

For sufficiently large values of n, i. e., for sufficiently small values of $\bar{x}$, $f'(\bar{x})$ is evidently negative. Hence for sufficiently large values of n we shall have

$$f'(1/2n\pi) < 0, \quad \text{and likewise} \quad f'(1/(2n+1)\pi) > 0.$$

Hence $f'(x)$ is both positive and negative at points on both sides of $x = 0$ inside of any interval about $x = 0$. And $f'(x)$ is continuous everywhere. *Nevertheless*, $f(x)$ is at an *absolute proper minimum* at $x = 0$.* For we have $f(0) = 0$, and $f(x) > 0$ when $x \neq 0$. Any attempt to establish the fact by any of the usual methods is futile, for $f''(x) = 0$ for $x = 0$ and $f'''(0)$ does not exist, while both f'' and f''' are both positive and negative near $x = 0$ on both sides of that point.

This example depends essentially, as do those which follow, upon the fact established above, viz., that $\xi(x, h)$ does not necessarily take on all values near zero for any limited values of h even though $f'(x)$ is continuous. For if ξ took on all values in the manner specified above, we should have

$$f'(k+t) > 0 \quad \text{and} \quad f'(k-t) < 0 \quad \text{whenever} \quad 0 < t < \delta,$$

since $f(k+h) - f(k) > 0$ and $f(k-h) - f(k) < 0$ provided $x = k$ gives a minimum. In fact we may say that

XX. *If the function* $|\xi(k, h)|$ *takes on all values less than* $^2\lambda$ *for some value of* $|h|$ *less than* $^1\rho$, *then the condition*

$$f'(k) = 0, \quad f'(k+t) > 0, \quad f'(k-t) < 0, \qquad 0 < t < \delta$$

is both necessary and sufficient for the existence of a proper minimum at $x = k$.

It is evident from the preceding example, however, that $f(x)$ may be at

* That is $f(0) < f(x)$ for all $x \neq 0$.

an absolute proper minimum and that $f'(x)$ may have no settled sign in any interval on either side of the point in question, even though $f'(x)$ is everywhere continuous.* Since the values which ξ takes on for values of h less than any positive number ρ form an assemblage with the power of the continuum, we may at least assert that

XXI. *If a continuous function $f(x)$ is at a proper minimum for $x = k$, the assemblage of values of x for which $f'(x)$ is negative (positive) has the power of the continuum in any interval whatever to the left (right) of $x = k$.*

In a similar manner examples may be constructed in which any desired number of derivatives exist and are continuous, but for which no process except direct inspection of the given function leads to any result; and for which the point in question gives an absolute proper minimum.

It may be objected that in each of the preceding examples the number of maxima and minima is infinite. In the following example:

$$y = \phi(x) = \int_0^x f(x)\,dx,$$

where

$$f(x) = x^3\left(1 + \sin\frac{1}{x}\right) \qquad \text{if } x \neq 0,$$

$$f(x) = 0 \qquad \text{if } x = 0,$$

this objection is removed. Since $f(x)$ is everywhere continuous and has the same sign as x itself, the function $\phi(x)$ is continuous, has its derivative $\phi'(x)\,[= f(x)]$ everywhere continuous, and is at an *isolated absolute proper minimum* for $x = 0$. *Nevertheless* $\phi'(x)$ does not always have the same sign as x, for $f(x)\,[= \phi'(x)]$ vanishes whenever $x = \dfrac{2}{(4n-1)\pi}$, and changes sign at such points.

While the results presented in this paper cannot be said to be complete, they present at least some characteristic properties of the remarkable function $\xi(x, h)$ which occurs in the law of the mean, and the writer hopes that this study may clear up some of the indistinct ideas regarding derivatives which must have troubled many a reader.

COLUMBIA, MISSOURI,
DECEMBER, 1905.

* That is, knowing that $f(x)$, together with $f'(x)$, is everywhere continuous, and knowing that $f(x)$ has a maximum for $x = c$, we can *not* assert that $f'(c + h) \leqq 0$ when $0 < h < \delta$, for example, even though δ be arbitrarily small. This holds even if the maximum is isolated and proper (see next example).

CONTENTS

PAGE

A Proof of the Theorem Concerning Artificial Singularities. By Professor D. R. Curtiss, ... 161

Another Proof of the Theorem Concerning Artificial Singularities. By Professor Maxime Bôcher, ... 163

Curves of Minimum Moment of Inertia with Respect to a Point. By Professor Max Mason, ... 165

Triangles and Quadrilaterals Inscribed to a Cubic and Circumscribed to a Conic. By Professor H. S. White, ... 172

On a Function which Occurs in the Law of the Mean. By Professor E. R. Hedrick, ... 177

ANNALS OF MATHEMATICS

Published in October, January, April, and July, under the auspices of Harvard University, Cambridge, Mass., U. S. A.

Cambridge: Address *The Annals of Mathematics*, 2 University Hall, Cambridge, Mass., U. S. A. Subscription price, $2 a volume (four numbers) in advance. Single numbers, 75c. All drafts and money orders should be made payable to Harvard University.

London: Longmans, Green & Co., 39 Paternoster Row. Price, 2 shillings a number.

Leipzig: Otto Harrassowitz, Querstrasse 14. Price, 2 marks a number.

Printed by The Salem Press Co., Salem, Mass., U. S. A.